arXiv:cond-mat/0001182v1 [cond-mat.supr-con] 13 Jan 2000

Unconventional strong pinning in the low temperature phase of $U_{.9725}Th_{.0275}Be_{13}$

Elisabeth Dumont [a,1], Ana Celia Mota [a], and James L. Smith [b]
[a] Laboratorium für Festkörperpkysik, ETH Hönggerberg, 8093 Zürich, Switzerland
[b] Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

Abstract

We investigated low field vortex dynamics in a single crystal of $U_{.9725}Th_{.0275}Be_{13}$.We found a sharp transition in the vortex creep rate at the lower transition temperature T_{c2}, coincident with the second jump in the specific heat. In the high-temperature phase, rather strong creep rates are observed. In the low temperature phase, the rates drop to undetectabely low levels. This behaviour indicates that a very strong pinning mechanism is present in the low temperature phase of $U_{.9725}Th_{.0275}Be_{13}$, which could be explained by the existence of domain walls, separating discreetly degenerate states of a superconductor, that can sustain fractional vortices and thus act as very strong pinning centers.

Key words: Heavy Fermions; Superconductivity; Vortex creep

If a small amount of uranium is substituted for thorium in the heavy fermion superconductor UBe_{13}, the superconducting transition temperature shows a non-monotonic dependence on thorium concentration x. Moreover, in a small critical range $0.019 < x < 0.045$ a second transition occurs below the superconducting transition[1]. The nature of this phase transition is still a matter of controversy. In zero field μSR measurements, a continuous increase of μ-spin-relaxation rate, which sets in at T_{c2} has been observed[2]. This has been interpreted as originating from a very weak spontaneous magnetic field. Superconducting origin of the transition at T_{c2} has been deduced from the observation of a distinct increase in the slope of the lower critical field H_{c1}[3] at T_{c2}.

Recently, the low-field magnetic properties of the heavy fermion superconductor UPt_3 have been investigated [4]. In the low–field, low–temperature B–phase, bulk vortices remain so strongly pinned that the creep rates drop from rather high finite rates observed in the A–phase to undetectabely low levels at T_c^-. The observation of zero initial creep in the B–phase of UPt_3 was then interpreted as originating from an intrinsic pinning mechanism in multicomponent superconducting phases. In such phases fractional vortices get trapped in domain walls separating discreetly degenerate states of the superconductor[5]. These experimental results show that vortex creep measurements can be

---

[1] Corresponding author. Present address: Laboratorium für Festkörperpkysik, ETH Hönggerberg, CH 8093 Zürich, Switzerland. E-mail: dumont@solid.phys.ethz.ch

Preprint submitted to Physica B *6 February 2008*

used as a powerful probe to give information about the character of a superconducting phase.

We present here relaxation measurements of the remanent magnetization on a single crystal of $U_{.9725}Th_{.0275}Be_{13}$(size: $1.00 \times 0.89 \times 2.25\,\mathrm{mm}^3$) prepared in Los Alamos National Laboratory[7]. The experimental arrangement has been described earlier [4]. The single crystal has a transition temperature $T_{c1} = 523\,\mathrm{mK}$. Specific heat measurements [6] show a second jump at $T_{c2} = 350\,\mathrm{mK}$. Typical relaxation measurements of the remanent magnetization are shown in fig. 1. All the values of M_{rem} are taken with the sample in the fully critical state.



Fig. 1. Isothermal relaxation measurements of remanent M_{rem} in $U_{.9725}Th_{.0275}Be_{13}$. Temperatures are given in mK on the right hand side of the graph.

We observe two different regimes of vortex creep, separated by a dramatic change in creep rates at $T = T_{c2}$. For temperatures above T_{c2}, we observe strong vortex creep and a logarithmic time dependence of the decays. For temperatures below T_{c2}, the initial vortex creep drops to zero. At longer times the decays start deviating from logarithmic behaviour. This acceleration of vortex creep at long times has also been observed in UPt_3[4]. Here we only discuss the initial slope of the decays. The normalized creep rates ($\partial \ln M/\partial \ln t$) are plotted in Fig.2 as a function of temperature. At $T = T_{c2}$, we observe a sharp transition in creep rates. They drop by 3 orders of magnitude to zero within our sensitivity ($\partial \ln M/\partial \ln t \approx 10^{-5}$). Also shown in Fig.2 are measurements of the lower critical field H_{c1} on the same crystal. H_{c1} was obtained as the field where the first deviation from the initial slope of isothermal magnetization–curves occurs.



Fig. 2. Creep rates (closed circles, right axis) and lower critical field H_{c1} (open diamonds, left axis) of $U_{.9725}Th_{.0275}Be_{13}$plotted vs T^2. The lines are guides to the eye.

In conclusion, we have observed a sharp transition in vortex pinning in $U_{.9725}Th_{.0275}Be_{13}$at T_{c2}, similar to the one observed in UPt_3 at T_c^-[4]. This might hint towards a new type of strong pinning structures and/or a new type of vortices being present in the low–T phases of these two multiphase superconductors.

References

[1] H. R. Ott et al., Phys. Rev. B **31** (1985) 1651.

[2] R. H. Heffner et al., Phys. Rev. B **40** (1989)806.

[3] U. Rauchschwalbe et al., Europhys. Lett. **3** (1987)757.

[4] A. Amann et al., Phys. Rev. B **57** (1998) 3640.

[5] M. Sigrist, K. Ueda, Rev. Mod. Phys. **63** (1992) 239.

[6] A. Ramirez, private communication.

[7] J.L. Smith, Phil. Mag. B, **65** (1992) 1367.